

HILBERT-SAMUEL POLYNOMIALS FOR THE CONTRAVARIANT EXTENSION FUNCTOR

ANDREW CRABBE, DANIEL KATZ, JANET STRIULI, AND EMANOIL THEODORESCU

ABSTRACT. Let (R, \mathfrak{m}) be a local ring and M and N finite R -modules. In this paper we give a formula for the degree of the polynomial giving the lengths of the modules $\text{Ext}_R^i(M, N/\mathfrak{m}^n N)$. A number of corollaries are given and more general filtrations are also considered.

1. INTRODUCTION

Let $(R, \mathfrak{m}, \mathfrak{k})$ be a Noetherian local ring, $I \subseteq R$ an ideal, M and N finitely generated R -modules. It is well known that if the lengths $\lambda(M/I^n M)$ of the modules $M/I^n M$ are finite for n large, these lengths are given by a rational polynomial of degree $\dim(M)$. In [7] (see also [6]) it is shown that the lengths of the modules $\text{Tor}_i^R(M, N/I^n N)$ and $\text{Ext}_R^i(M, N/I^n N)$ have polynomial growth for large n , whenever the lengths of these modules are finite. However, the degrees of the corresponding Hilbert-Samuel polynomials are not as easy to determine (see [7], [3], and [5]). The purposes of this note are three-fold. The first is to improve the known estimates for the degrees of the polynomials giving the lengths of $\text{Ext}_R^i(M, N/I^n N)$ and $\text{Tor}_i^R(M, N/I^n N)$ in the case $I = \mathfrak{m}$ by giving a precise formula for these degrees. Previous results for the case $I = \mathfrak{m}$ for the torsion functor were given in [3] and [5], where various assumptions were made in order to control this degree. In this paper we do not need to make any assumptions on M , N or R to obtain our formulas, and we need only make modest assumptions on them to obtain a formula that makes direct reference only to M and N . In fact, in section two we begin by giving a general formula (see Proposition 2.1) for the degree of the Hilbert polynomial associated to general cohomology or homology modules, which specializes to our main results in section three when the ideal in question is \mathfrak{m} and the cohomology is determined by either the contravariant extension functor (Theorem 3.2) or the torsion functor (Theorem 3.12). For example, the following theorem is a direct consequence of our main result in section three. Here we write $\mathcal{E}_{M,N,\mathfrak{m}}^i(n)$ for the Hilbert polynomial giving $\lambda(\text{Ext}_R^i(N, M/\mathfrak{m}^n M))$, for n large.

Theorem 1.1. *Let $(R, \mathfrak{m}, \mathfrak{k})$ be a local Noetherian ring and let M and N be two finitely generated R -modules such that M has a rank and $\dim(N) = \dim(R)$. Then $\deg(\mathcal{E}_{M,N,\mathfrak{m}}^i(n)) = \dim(N) - 1$.*

As we will see below, the degree of $\mathcal{E}_{M,N,\mathfrak{m}}^i(n)$ is partially controlled by the dimension of $\Omega_R^i(M)$, the i^{th} syzygy of M . Consequently, as an application of our

2000 *Mathematics Subject Classification.* Primary 13D07, 13D40, 13H15, 13C11.

Key words and phrases. Hilbert-Samuel polynomial, extension functor, injective resolution.

The third author was partially supported by NSF grant DMS-0201904.

degree formula we obtain the following proposition, which yields some information about the dimension of the syzygies of finite length modules.

Proposition 1.2. *Let $(R, \mathfrak{m}, \mathfrak{k})$ be a local ring and M a finitely generated R -module, free of constant rank on the punctured spectrum of R . Assume $\dim(R) \geq 2$, that the betti numbers of M are non-decreasing and that $i < \text{p.d.}(M)$. Then $\dim(\Omega_R^{i+1}(M)) = d$.*

Our second purpose, especially in regards to section three, is to lay the groundwork for results concerning indecomposable modules in [2], where knowledge of the relative growth of the Hilbert polynomials of large syzygies of the residue field \mathfrak{k} is required. In particular, Theorem 1.1 above plays a crucial role in [2].

Finally, in section four, we address our third purpose where our goal is to show that the results of section three can, in many cases, be extended to more general filtrations to give results for the extension functor parallel to those given in [5] for the torsion functor.

2. GENERAL COHOMOLOGY

Throughout, $(R, \mathfrak{m}, \mathfrak{k})$ will denote a Noetherian local ring of Krull dimension d and all modules will be finitely generated R -modules. In this section we prove a general result about the Hilbert polynomial associated to an ideal and the cohomology (or homology) of a complex. We start by letting

$$\mathcal{C} : \quad X \xrightarrow{\alpha} Y \xrightarrow{\beta} Z$$

be a complex of finitely generated R -modules with $Y \neq 0$. We assume that $I \subseteq R$ is an ideal such that the homology modules $H(\mathcal{C} \otimes R/I^n)$ associated to $\mathcal{C} \otimes R/I^n$ have finite length for n large. By [7], Proposition 3, the lengths of the modules $H(\mathcal{C} \otimes R/I^n)$ are given by a polynomial $P_I^{\mathcal{C}}(n)$ for n large. The following proposition strengthens [7], Proposition 3, in that we replace the degree estimate there by an equality. In the statement of this proposition we set

$$\mathcal{M} := \bigoplus_{n \geq 0} (I^n Z \cap \text{im}(\beta)) / I^n \text{im}(\beta).$$

Note that \mathcal{M} is a finitely generated graded module over the Rees algebra of R with respect to I , so that if its graded components have finite length as R -modules, then these lengths are ultimately given by a rational polynomial of degree $\dim(\mathcal{M}) - 1$.

Proposition 2.1. *Let $(R, \mathfrak{m}, \mathfrak{k})$ be a local ring and \mathcal{C} as above be a complex of finitely generated R -modules with $Y \neq 0$. Let $I \subseteq R$ be an ideal such that the lengths of the cohomology modules $H(\mathcal{C} \otimes R/I^n)$ are finite for n large and let $P_I^{\mathcal{C}}(n)$ denote the corresponding Hilbert-Samuel polynomial. Then*

$$\deg(P_I^{\mathcal{C}}(n)) = \max\{\dim(H(\mathcal{C})), \dim(\mathcal{M}) - 1\}.$$

Proof. Set $A := \ker(\beta)$ and $B := \text{im}(\alpha)$. We begin by arguing as in the proof of [7], Proposition 3. By the Artin-Rees Lemma, there exists $h > 0$ so that for $n \geq h$,

$$I^n Z \cap \text{im}(\beta) = I^{n-h}(I^h Z \cap \text{im}(\beta)).$$

Since an element in the cohomology of the complex $\mathcal{C} \otimes R/I^n$ corresponds to an element in Y that gets mapped by β into $I^n Z \cap \text{im}(\beta)$, it follows (see [7]) that for

$n \geq h$,

$$\mathbf{H}(\mathcal{C} \otimes R/I^n) = \frac{A + I^{n-h}C}{B + I^{n-h}D},$$

where $C := \beta^{-1}(I^h Z)$ and $D := I^h Y$. Now for n large,

$$P_I^{\mathcal{C}}(n) = \lambda\left(\frac{A + I^{n-h}C}{B + I^{n-h}D}\right) = \lambda\left(\frac{A + I^{n-h}C}{A + I^{n-h}D}\right) + \lambda\left(\frac{A + I^{n-h}D}{B + I^{n-h}D}\right) \quad (2.1)$$

$$= \lambda\left(\frac{A + I^{n-h}C}{A + I^{n-h}D}\right) + \lambda\left(\frac{U + I^{n-h}W}{I^{n-h}W}\right), \quad (2.2)$$

where $U := A/B = \mathbf{H}(\mathcal{C})$ and $W := (D + B)/B$. We first note that by [7], Lemma 2, both length expressions on the right hand sides of the displayed equations (2.1) and (2.2) are given by polynomials. Let $P_1(n)$ denote the polynomial giving the lengths of $(A + I^{n-h}C)/(A + I^{n-h}D)$ and $P_2(n)$ denote the polynomial giving the lengths of $(U + I^{n-h}W)/I^{n-h}W$. We first calculate the degree of $P_1(n)$. For this, we note that, by definition,

$$\frac{A + I^{n-h}C}{A + I^{n-h}D} \cong \frac{I^n Z \cap \text{im}(\beta)}{I^n \text{im}(\beta)}. \quad (2.3)$$

Thus, by the definition of \mathcal{M} , $\deg(P_1(n)) = \dim(\mathcal{M}) - 1$.

We now show that the degree of $P_2(n)$ equals $\dim(U) = \dim(\mathbf{H}(\mathcal{C}))$. Since $P_I^{\mathcal{C}}(n) = P_1(n) + P_2(n)$, this will complete the proof of the proposition. For this note that there exists $c > 0$ so that for n sufficiently large,

$$P_2(n) = \lambda\left(\frac{U}{U \cap I^{n-h}W}\right) = \lambda\left(\frac{U}{U \cap I^c W}\right) + \lambda\left(\frac{U \cap I^c W}{I^{n-h-c}(U \cap I^c W)}\right),$$

so $P_2(n)$ has degree equal to $\dim(U \cap I^c W)$. But since $U/(U \cap I^c W)$ has finite length, $\dim(U) = \dim(U \cap I^c W)$, which gives what we want. \square

In the case that $I = \mathfrak{m}$ we can replace \mathcal{M} in the statement of Proposition 2.1 by $T := \text{im}(\beta)$. This allows us to give a precise formula for $P_{\mathfrak{m}}^{\mathcal{C}}(n)$ in terms of the modules appearing in the complex \mathcal{C} . Our main results in the next section concerning the extension and torsion functors are immediate consequences of the following theorem.

Theorem 2.2. *Let $(R, \mathfrak{m}, \mathfrak{k})$ be a local ring and \mathcal{C} as above be a complex of finitely generated R -modules with $Y \neq 0$. Write $P_{\mathfrak{m}}^{\mathcal{C}}(n)$ for the Hilbert-Samuel polynomial giving the lengths of the cohomology modules $\mathbf{H}(\mathcal{C} \otimes R/\mathfrak{m}^n)$, for n large and set $T := \text{im}(\beta)$. If $T \subseteq \mathfrak{m}Z$, then*

$$\deg(P_{\mathfrak{m}}^{\mathcal{C}}(n)) = \max\{\dim(\mathbf{H}(\mathcal{C})), \dim(T) - 1\}.$$

Proof. By Proposition 2.1, we just have to show that $\dim(\mathcal{M}) = \dim(T)$. On the one hand, since, $T \subseteq \mathfrak{m}Z$, we have

$$\mathfrak{m}^{n-1}T \subseteq \mathfrak{m}^n Z \cap T.$$

Therefore, $\bigoplus_{n \geq 1} \mathfrak{m}^{n-1}T/\mathfrak{m}^n T \subseteq \mathcal{M}$. It follows that

$$\dim(T) = \dim\left(\bigoplus_{n \geq 1} \mathfrak{m}^{n-1}T/\mathfrak{m}^n T\right) \leq \dim(\mathcal{M}).$$

On the other hand, by the Artin-Rees lemma, there exists an $e \geq 1$ so that for n large, $\mathfrak{m}^n Z \cap T = \mathfrak{m}^{n-e}(\mathfrak{m}^e Z \cap T)$. As in the proof of Proposition 2.1, write $P_1(n)$ for the polynomial giving the lengths of \mathcal{M}_n . Then

$$P_1(n) \leq \lambda(T/\mathfrak{m}^n T) - \lambda(T/\mathfrak{m}^{n-e} T),$$

and the degree of the polynomial giving the latter difference equals $\dim(T) - 1$. This shows that \mathcal{M} has dimension less than or equal to the dimension of T and thus must have dimension equal to T , which is what we want. \square

In the following corollary, we record some observations related to the case that $P_I^{\mathcal{C}}(n)$ is the zero polynomial.

Corollary 2.3. *Retain the notation from Proposition 2.1 and Theorem 2.2.*

- (a) *If $P_I^{\mathcal{C}}(n) \equiv 0$, then $H(\mathcal{C}) = 0$.*
- (b) *If $\text{im}(\alpha) \subseteq \mathfrak{m}Y$, then $H(\mathcal{C})$ and $\text{im}(\beta)$ cannot be simultaneously zero.*
- (c) *If $\text{im}(\alpha) \subseteq \mathfrak{m}Y$ and $\text{im}(\beta) \subseteq \mathfrak{m}Z$, then $P_{\mathfrak{m}}^{\mathcal{C}}(n) \equiv 0$ if and only if $H(\mathcal{C}) = 0$ and $\text{im}(\beta)$ has non-zero finite length.*

Proof. Part (a) is immediate from equation (2.2) in the proof Proposition 2.1. For part (b), suppose $\text{im}(\beta) = 0$. Then $H(\mathcal{C}) = Y/\text{im}(\alpha)$. Since $\text{im}(\alpha) \subseteq \mathfrak{m}Y$ and $Y \neq 0$, we cannot have $H(\mathcal{C}) = 0$, by Nakayama's lemma.

For part (c), suppose first that $P_{\mathfrak{m}}^{\mathcal{C}}(n) \equiv 0$. By part (a), we have $H(\mathcal{C}) = 0$. From equations (2.1) and (2.3) we have $\mathfrak{m}^n Z \cap \text{im}(\beta) = \mathfrak{m}^n \text{im}(\beta)$, for n large. Since $\text{im}(\beta)$ is contained in $\mathfrak{m}Z$, it follows that $\mathfrak{m}^{n-1} \text{im}(\beta) = \mathfrak{m}^n \text{im}(\beta)$, so by Nakayama's lemma, $\text{im}(\beta)$ has finite length. By part (b), $\text{im}(\beta) \neq 0$. Conversely, suppose that $H(\mathcal{C}) = 0$ and $\text{im}(\beta)$ has finite length. Using the notation from the proof of Proposition 2.1, the fact that $U := H(\mathcal{C}) = 0$ implies $P_2(n) \equiv 0$. Since $\text{im}(\beta)$ has finite length, equation (2.3) and the Artin-Rees lemma imply $P_1(n) \equiv 0$. Since $P_{\mathfrak{m}}^{\mathcal{C}}(n) = P_1(n) + P_2(n)$, this gives what we want. \square

Remark 2.4. In regards to Proposition 2.1, we would like to point out that generally, it is the term $\dim(\mathcal{M}) - 1$ that makes determining the exact degree of $P_I^{\mathcal{C}}(n)$ for arbitrary I difficult. Indeed, whenever $H(\mathcal{C})$ is zero, then the degree of $P_I^{\mathcal{C}}(n)$ is equal to $\dim(\mathcal{M}) - 1$. (Think of the case when I is \mathfrak{m} -primary and $P_I^{\mathcal{C}}(n) = \lambda(\text{Tor}_i(R/I^n, R))$.) Our success in determining the degree of $P_{\mathfrak{m}}^{\mathcal{C}}(n)$ in Theorem 2.2 and the corresponding results in section three is due to the fact that in these cases, we can calculate the dimension of \mathcal{M} . Similarly, our success in section four with filtrations more general than the \mathfrak{m} -adic filtration is due strictly to the ability to calculate the dimension of \mathcal{M} in those cases as well.

3. THE \mathfrak{m} -ADIC FILTRATION FOR THE CONTRAVARIANT EXTENSION FUNCTOR

In this section we apply the results of the previous section to give a precise formula for the degree of the Hilbert polynomial giving the lengths of the modules $\text{Ext}_R^i(M, N/\mathfrak{m}^n N)$ and $\text{Tor}_i^R(M, N/\mathfrak{m}^n N)$. Our formulas below for the degrees of these polynomials (see Theorems 3.2 and 3.12) involve the dimension of the image of the i^{th} syzygy of M or its transpose in an appropriate direct sum of copies of N , and are immediate consequences of Theorem 2.2. We will see that in those cases where the dimension of syzygies of M and their transposes are well-behaved, our formulas either agree with or improve prior estimates. However, the formulas in Theorems 3.2 and 3.12 are valid in all cases.

We begin by establishing some notation. Let

$$\mathbf{F} : \quad \cdots \rightarrow F_{i+1} \xrightarrow{\phi_{i+1}} F_i \xrightarrow{\phi_i} F_{i-1} \rightarrow \cdots$$

denote a minimal free resolution of M . We set $\beta_i(M) := \text{rank}(F_i)$ for all i . Thus, for all $i \geq 0$, $\beta_i(M)$ is the i^{th} betti number of M . We can calculate $\text{Ext}^i(M, N/I^n N)$ by applying $\text{Hom}(-, N/I^n N)$ to \mathbf{F} . Thus, $\text{Ext}^i(M, N/I^n N)$ is the cohomology of the co-chain complex $\text{Hom}(\mathbf{F}, N/I^n N)$.

Alternately, we may first apply the functor $\text{Hom}(-, N)$ to the resolution \mathbf{F} to obtain the complex

$$\text{Hom}(\mathbf{F}, N) : \quad \cdots \rightarrow N^{\beta_{i-1}(M)} \xrightarrow{\overline{\phi_i}} N^{\beta_i(M)} \xrightarrow{\overline{\phi_{i+1}}} N^{\beta_{i+1}(M)} \rightarrow \cdots,$$

which we then tensor with R/I^n . The resulting complex $\text{Hom}_R(\mathbf{F}, N) \otimes R/I^n$ is isomorphic to $\text{Hom}_R(\mathbf{F}, N/I^n N)$. Hence we may calculate $\text{Ext}_R^i(M, N/I^n N)$ as the i^{th} cohomology of $\text{Hom}_R(\mathbf{F}, N) \otimes R/I^n$. Homology and cohomology of complexes of this form were studied in [7].

Hilbert polynomials for derived functors. For fixed $i \geq 1$, assume that the modules $\text{Ext}^i(M, N/I^n N)$ have finite length for n large. It follows from [7], Corollary 4, that the lengths $\lambda(\text{Ext}_R^i(M, N/I^n N))$ are given by a rational polynomial for n sufficiently large. We will write $\mathcal{E}_{M,N,I}^i(n)$ for this polynomial. By Corollary 4 in [7], we have the following estimate for the degree of $\mathcal{E}_{M,N,I}^i(n)$.

$$\deg(\mathcal{E}_{M,N,I}^i(n)) \leq \max\{\dim(\text{Ext}_R^i(M, N)), \ell_N(I) - 1\},$$

where $\ell_N(I)$ denote the analytic spread of I on N . Recall that $\ell_N(I)$ is the Krull dimension of the graded module $\bigoplus_{n \geq 0} I^n N / \mathfrak{m} I^n N$. Equivalently, if we write S for the ring $R / \text{ann}(N)$ and $\mathcal{R}(IS)$ for the Rees ring of S with respect to IS , then $\ell_N(I)$ is the dimension of the ring $\mathcal{R}(IS) / \mathfrak{m} \mathcal{R}(IS)$, i.e., it is just the analytic spread of the image of I in the ring S . (See [7], the proof of Proposition 3, for a proof of this fact.) Moreover, in [7] it is shown that equality holds in the degree estimate when the first term on the right is at least as large as the second term on the right. Similarly, in [7], it is shown that when, for $i \geq 1$ fixed, the modules $\text{Tor}_i^R(M, N/I^n N)$ have finite length, those lengths are given by a rational polynomial in n , for n sufficiently large. We will write $\tau_i^{M,N,I}(n)$ for the corresponding polynomial. In [7], it is shown that

$$\deg(\tau_i^{M,N,I}(n)) \leq \max\{\dim(\text{Tor}_i(M, N)), \ell_N(I) - 1\},$$

and equality holds when the first term on the right hand side of the inequality is at least as large as the second term on the right.

Remark 3.1. Before stating one of the main results of this section, we first point out that the degree bounds above can be recovered from Proposition 2.1. For the degree of $\mathcal{E}_{M,N,I}^i(n)$, it follows from Proposition 2.1 that

$$\deg(\mathcal{E}_{M,N,I}^i(n)) = \max\{\dim(\text{Ext}_R^i(M, N)), \dim(\mathcal{T}_i) - 1\},$$

for $\mathcal{T}_i = \bigoplus_{n \geq 0} (I^n N^{\beta_{i+1}(M)} \cap \text{im}(\overline{\phi_{i+1}})) / I^n \text{im}(\overline{\phi_{i+1}})$. Note that we can regard \mathcal{T}_i as a module over the Rees ring $\mathcal{R}(IS)$ of $S := R / \text{ann}(N)$ with respect to IS . Since, by assumption, the lengths of the graded components of \mathcal{T}_i are finite (see the proof of Proposition 2.1), \mathcal{T}_i must be annihilated by some power of $\mathfrak{m} \mathcal{R}(IS)$, say $\mathfrak{m}^q \mathcal{R}(IS)$. Then $\dim(\mathcal{T}_i) \leq \dim(\mathcal{R}(IS) / \mathfrak{m}^q \mathcal{R}(IS)) = \ell_N(I)$, which gives what we want. The argument for the degree bound involving $\tau_i^{M,N,I}(n)$ is entirely analogous.

We are now ready for the main result of this section.

Theorem 3.2. *Let (R, \mathfrak{m}) be a local ring, M and N finitely generated R -modules. Fix $0 \leq i \leq \text{p.d.}(M)$. Set $T_i := \text{im}(\overline{\phi_{i+1}})$, for $\overline{\phi_{i+1}}$ as above. Then*

$$\deg(\mathcal{E}_{M,N,\mathfrak{m}}^i(n)) = \max\{\dim(\text{Ext}^i(M, N)), \dim(T_i) - 1\}.$$

Proof. If we use the fact above that $\text{Ext}_R^i(M, N/\mathfrak{m}^n N)$ is the i^{th} cohomology of the complex $\text{Hom}(\mathbf{F}, N) \otimes R/\mathfrak{m}^n$, then the theorem follows immediately from Theorem 2.2. \square

Remark 3.3. (i) Since the resolution \mathbf{F} is minimal, it follows from Corollary 2.3 that $\mathcal{E}_{M,N,\mathfrak{m}}^i(n)$ is identically zero if and only if $\text{Ext}^i(M, N) = 0$ and T_i has non-zero finite length. Therefore, if we adopt the conventions that the zero polynomial has degree -1 and the zero module has dimension $-\infty$, Theorem 3.2 does give the correct value for the degree of $\mathcal{E}_{M,N,\mathfrak{m}}^i(n)$ in this case.

(ii) Concerning the upper bound for the degree of $\mathcal{E}_{M,N,I}^i(n)$ given before Remark 3.1, when I is an \mathfrak{m} -primary ideal, in particular, when $I = \mathfrak{m}$, $\ell_N(I) = \dim(N)$. Thus, since $\dim(T_i) \leq \dim(N)$, we see that Theorem 3.2 improves the estimate from [7] in the special case that $\mathfrak{m} = I$. This improvement is extended to more general filtrations (but not all modules) in the next section.

(iii) The Hilbert polynomial $\tau_i^{M,R,\mathfrak{m}}(n)$ giving the lengths of $\text{Tor}_i(M, R/\mathfrak{m}^n)$ has degree less than d for all $i > 0$ (see [5] or Theorem 3.12 below). This no longer holds for $\mathcal{E}_{M,R,\mathfrak{m}}^i(n)$. Indeed, let R be a local ring with a prime ideal P of maximal dimension such that R_P is not Gorenstein and set $M := R/P$. Then $\text{Ext}^i(M, R)_P \neq 0$ for all $i > 0$, so that $\dim(\text{Ext}^i(M, R)) = d$, for all $i > 0$. Thus, by Theorem 3.2, $\deg(\mathcal{E}_{M,R,\mathfrak{m}}^i(n)) = d$, for all $i > 0$.

Corollary 3.4. *Suppose that M has finite projective dimension, say $\text{p.d.}(M) = i$. Then $\deg(\mathcal{E}_{M,N,\mathfrak{m}}^i(n)) = \dim(\text{Ext}^i(M, N))$.*

Proof. This is immediate from the theorem, since in this case, $T_i = 0$. \square

In Theorem 3.2, we may replace T_i by N in any number of situations as the following corollary shows.

Corollary 3.5. *Let (R, \mathfrak{m}) be a local ring, M and N finitely generated R -modules. Fix $0 \leq i < \text{p.d.}(M)$ and suppose that one of the following conditions holds.*

- (a) $i = 0$.
- (b) *There exists a prime ideal P of maximal dimension in the support of N so that some entry of ϕ_{i+1} , say r , does not belong to P (e.g., r is a non-zero divisor on N).*
- (c) $\beta_i(M) > \beta_{i-1}(M)$.
- (d) $N = R$ and M_P is not a free R_P -module, for some prime ideal $P \subseteq R$ of dimension d .

Then

$$\deg(\mathcal{E}_{M,N,\mathfrak{m}}^i(n)) = \max\{\dim(\text{Ext}^i(M, N)), \dim(N) - 1\}.$$

Proof. For (a), we have an exact sequence

$$0 \rightarrow \text{Hom}(M, N) \rightarrow N^{\beta_0} \xrightarrow{\overline{\phi_1}} T_0 \rightarrow 0.$$

By Theorem 3.2, if $\dim(\text{Hom}(M, N)) = \dim(N)$, $\deg(\mathcal{E}_{M,N,I}^0(n)) = \dim(N)$, which is the maximum value in question. Otherwise, the exact sequence above shows that T_0 and N have the same dimension, so $\dim(T_0) - 1 = \dim(N) - 1$ is the maximum value, and this gives what we want.

For (b), since $r_P \cdot N_P \neq 0$, $(T_i)_P = \text{im}(\overline{\phi_{i+1}})_P \neq 0$. Thus, $\dim(T_i) = \dim(N)$, and this gives what we want, by Theorem 3.2.

To prove (c), let P be a prime ideal of maximal dimension in the support of N . By Theorem 3.2, it suffices to show that either $\text{Ext}^i(M, N)_P \neq 0$ or $(T_i)_P \neq 0$. If $\text{Ext}^i(M, N)_P = 0$ and $(T_i)_P = 0$, then $(\overline{\phi_i})_P$ is surjective. But this cannot happen if $\beta_i(M) > \beta_{i-1}(M)$.

For (d), by assumption, $(\phi_{i+1})_P \neq 0$, so $(T_i)_P \neq 0$, which gives the result. \square

It is clear that as long as any one of the conditions (a)-(d) of the previous corollary is met, we obtain $\deg(\mathcal{E}_{M,N,m}^i(n)) = \dim(N) - 1$, whenever the dimension of $\text{Ext}^i(M, N)$ is less than $\dim(N)$. We list a couple of such cases in the following corollary.

Corollary 3.6. *Let (R, \mathfrak{m}) be a local ring, M and N finitely generated R -modules. Fix $0 \leq i < \text{p.d.}(M)$ and suppose that one of the following conditions holds.*

- (a) M has a rank and $\dim(N) = \dim(R)$.
- (b) M is a non-zero syzygy of \mathfrak{k} and $\dim(N) \geq 1$.

Then $\deg(\mathcal{E}_{M,N,m}^i(n)) = \dim(N) - 1$.

Proof. Let P be a prime of maximal dimension in the support of N . In case (a), P is a minimal prime of R . Therefore, by [5], Remark 2.1, in either case (a) or case (b), the image of $(\phi_{i+1})_P$ is a non-zero summand of $(F_i)_P$. Thus, in each case, at least one entry of $(\phi_{i+1})_P$ is a unit. Therefore, in both cases, the conclusion of Corollary 3.5 holds, i.e.,

$$\deg(\mathcal{E}_{M,N,m}^i(n)) = \max\{\dim(\text{Ext}^i(M, N)), \dim(N) - 1\}.$$

On the other hand, if (a) holds, it follows that $\dim(\text{Ext}^i(M, N)) < d = \dim(N)$, so the maximum value in question is $\dim(N) - 1$. Similarly, if hypothesis (b) holds, then $\dim(\text{Ext}^i(M, N)) = 0$, so again the maximum in question is $\dim(N) - 1$, which completes the proof. \square

Corollary 3.7. *If $\beta_i(M) = \beta_{i-1}(M)$, then $\deg(\mathcal{E}_{M,N,m}^i(n)) \geq \dim(N) - 2$.*

Proof. By dimension shifting, we may assume $i = 1$. Then we have an exact sequence

$$0 \rightarrow \frac{\mathfrak{m}^n N}{\mathfrak{m}^{n+1} N} \rightarrow \frac{N}{\mathfrak{m}^{n+1} N} \rightarrow \frac{N}{\mathfrak{m}^n N} \rightarrow 0,$$

from which we obtain

$$\begin{aligned} 0 \rightarrow \text{Hom}(M, \mathfrak{m}^n N / \mathfrak{m}^{n+1} N) &\rightarrow \text{Hom}(M, N / \mathfrak{m}^{n+1} N) \rightarrow \text{Hom}(M, N / \mathfrak{m}^n N) \\ &\rightarrow \text{Ext}^1(M, \mathfrak{m}^n N / \mathfrak{m}^{n+1} N) \xrightarrow{\rho_n} \text{Ext}^1(M, N / \mathfrak{m}^{n+1} N). \end{aligned} \quad (3.1)$$

Since the resolution of M is minimal the lengths of

$$\text{Hom}(M, \mathfrak{m}^n N / \mathfrak{m}^{n+1} N) \quad \text{and} \quad \text{Ext}^1(M, \mathfrak{m}^n N / \mathfrak{m}^{n+1} N)$$

are just $\beta_0(M) \cdot L(n)$ and $\beta_1(M) \cdot L(n)$ respectively, where $L(n)$ is the polynomial giving the minimal number of generators of $\mathfrak{m}^n N$ for n large. Using the fact that $\beta_1(M) = \beta_0(M)$, it follows that

$$\mathcal{E}_{M,N,\mathfrak{m}}^1(n+1) \geq \mathcal{E}_{M,N,\mathfrak{m}}^0(n+1) - \mathcal{E}_{M,N,\mathfrak{m}}^0(n).$$

By Corollary 3.5, $\mathcal{E}_{M,N,\mathfrak{m}}^0(n)$ has degree greater than or equal to $\dim(N) - 1$, and this gives what we want. \square

Remark 3.8. Let R be a two dimensional local ring of depth one and suppose $a \in R$ is a parameter such that $(0 : (0 : a)) = (a)$. Then for $M := R/(a)$ and $i = 1$, we have that $\mathcal{E}_{M,R,\mathfrak{m}}^1(n)$ is a non-zero constant. In particular, this gives an example where $\beta_0(M) = \beta_1(M)$ and $\deg(\mathcal{E}_{M,R,\mathfrak{m}}^1(n)) = \dim(R) - 2$, i.e., an example where the lower bound in Corollary 3.7 is attained. To see this, let

$$\cdots \rightarrow R^r \xrightarrow{\phi_2} R \xrightarrow{a} R \rightarrow M \rightarrow 0$$

be the start of a minimal resolution of M . Note that the image of ϕ_2 is just $(0 : a)$. To calculate $\text{Ext}^1(M, R)$, we look at the dual of the resolution, thereby getting the complex

$$0 \rightarrow R \xrightarrow{a} R \xrightarrow{\phi_2^t} R^r \rightarrow \cdots$$

Note that $\ker(\phi_2^t) = (0 : (0 : a))$, so by our assumption, $\text{Ext}^1(M, R) = 0$. On the other hand, clearly a belongs to the annihilator of $T := \text{im}(\phi_2^t)$, so $\dim(T) \leq 1$. Since the depth of R is one, T cannot be zero-dimensional. Thus, $\dim(T) = 1$, so by Theorem 3.2, $\mathcal{E}_{M,R,\mathfrak{m}}^1(n)$ is a non-zero constant.

To find a concrete example with the stated properties, it suffices to find a one dimensional local ring (S, \mathfrak{n}) with a parameter a satisfying $(0 : (0 : aS)) = (a)$. Indeed, given such an S and a , let $R := S[X]_{(\mathfrak{n}, X)}$, where X is an indeterminate over S . Then R and a meet the requirements stated above. Finally, to find such an S , we use the following example shown to us by Craig Huneke.

Example 3.9. Let k be a field and x, y, z, u, v be indeterminates. Let S denote the power series ring $k[[x, y, z, u, v]]$ modulo the ideal I , where I is the ideal generated by $x^2, xz, z^2, xu, zv, u^2, v^2, zu + xv + uv, yu, yv, yx - zu, yz - xv$. Then S is a one dimensional local ring with parameter ideal yS satisfying $(0 : (0 : yS)) = yS$.

For a deeper analysis of this situation, the interested reader should consult [4].

The previous two corollaries together with Theorem 3.2 yield some information on the dimension of syzygies over an arbitrary local ring. In the appendix we record some of the consequences our work has for the dimension of syzygies.

In our last corollary of this section concerning the contravariant extension functor, we record what happens when $N = R$ is a Cohen-Macaulay local ring with a canonical module.

Corollary 3.10. *Suppose that R is a Cohen-Macaulay local ring with canonical module ω . Assume $0 \leq i < \text{p.d.}(\omega)$. Then,*

- (a) $\deg(\mathcal{E}_{\omega,R,\mathfrak{m}}^i(n)) = \max\{\dim(\text{Ext}^i(\omega, R)), \dim(R) - 1\}$.
- (b) $\deg(\mathcal{E}_{\omega,R,\mathfrak{m}}^i(n)) = \dim(R) - 1$, if R is generically Gorenstein.

Moreover, if $i = 0$, then $\deg(\mathcal{E}_{\omega,R,\mathfrak{m}}^0(n)) = \dim(R)$.

Proof. We consider (a) and (b) together. Suppose there exists a minimal prime $P \subseteq R$ such that R_P is not Gorenstein. Then ω_P is not a free R -module, so (a) holds by Corollary 3.5d. Otherwise, R is generically Gorenstein, so (b) holds by Corollary 3.6a.

Now suppose $i = 0$ and let $P \subseteq R$ be a prime of maximal dimension. Then $\text{Hom}(\omega, R)_P \neq 0$, so $\dim(\text{Hom}(\omega, R)) = d$, and the result follows from part (a). \square

Remark 3.11. Returning to the set up at the beginning of this section, let $\tau_i^{M,N,\mathfrak{m}}(n)$ denote the polynomial giving the lengths of $\text{Tor}_i(M, N/\mathfrak{m}^n N)$, for n large. Then $\text{Tor}_i(M, N/\mathfrak{m}^n N)$ is the i^{th} homology in the complex $(\mathcal{F} \otimes N) \otimes R/\mathfrak{m}^n$, so that Theorem 2.2 yields a formula for the degree of $\tau_i^{M,N,\mathfrak{m}}(n)$ analogous to the one obtained in Theorem 3.2 for $\mathcal{E}_{M,N,\mathfrak{m}}^i(n)$. When $N = R$, several special cases were given in [5]. Note that one can then list several of corollaries to Theorem 3.12 analogous to those above.

Theorem 3.12. *Let (R, \mathfrak{m}) be a local ring and M and N finite R -modules. Fix $0 \leq i \leq \text{p.d.}(M)$ and let $\tau_i^{M,N,\mathfrak{m}}(n)$ denote the Hilbert polynomial giving the lengths of the modules $\text{Tor}_i(M, N/\mathfrak{m}^n N)$ for n large. Let C_i denote the image of the induced map $\bar{\phi}_i \otimes \bar{1}_N : F_i \otimes N \rightarrow F_{i-1} \otimes N$. Then*

$$\deg(\tau_i^{M,N,\mathfrak{m}}(n)) = \max\{\dim(\text{Tor}_i(M, N)), \dim(C_i) - 1\}.$$

In particular, if $N = R$ and $i \geq 1$, then

$$\deg(\tau_i^{M,N,\mathfrak{m}}(n)) = \dim(\Omega_R^i(M)) - 1.$$

Proof. For the first statement, we just apply Theorem 2.2 to the i^{th} spot of the complex $\mathbf{F} \otimes N$, where as before, \mathbf{F} denotes the minimal resolution of M . The second statement follows immediately from the first. \square

Remark 3.13. (i) Assume $N = R$ and $i \geq 1$. Then the theorem above shows that the degree of $\tau_i^{M,N,\mathfrak{m}}(n)$ is simply the dimension of the i^{th} syzygy of M minus one. On the other hand, for $N = R$ and $i \geq 1$, the degree of $\mathcal{E}_{M,N,\mathfrak{m}}^i(n)$ is determined by both the dimension of the module $\text{Ext}^i(M, R)$ and the dimension of T_i , which in this case is the dimension of the $(i+1)^{\text{st}}$ syzygy of M . Suppose M and R are such that $\text{Ext}^i(M, R)$ has dimension less than or equal to $d-1$ (e.g., R is generically Gorenstein or M has a rank). Then the degree of $\mathcal{E}_{M,R,\mathfrak{m}}^i(n)$ is less than or equal to $d-1$. Now, since either the i^{th} or the $(i+1)^{\text{st}}$ syzygy of M must have dimension equal to the dimension of R , it follows that either $\tau_i^{M,R,\mathfrak{m}}(n)$ or $\mathcal{E}_{M,R,\mathfrak{m}}^i(n)$ has maximal degree $\dim(R) - 1$. Of course, as above, very minor assumptions on M will also guarantee that both polynomials have maximal degree. However, as pointed out in Question 5.1, for R and M arbitrary, it is not known whether the dimensions of the syzygies of M ultimately stabilize, so one cannot make a definitive statement regarding the degrees of $\mathcal{E}_{M,R,\mathfrak{m}}^i(n)$ and $\tau_i^{M,R,\mathfrak{m}}(n)$, even for i sufficiently large. (ii) In [3], it is shown that for $i > 0$,

$$\text{depth}(R) - 1 \leq \deg(\tau_i^{M,R,\mathfrak{m}}(n)) \leq \dim(R) - 1.$$

We get by Theorem 3.12 that $\deg(\tau_i^{M,R,\mathfrak{m}}(n)) = \dim(\Omega_R^i(M)) - 1$. Now, on the one hand $\text{depth}(R) = \text{depth}(F_i) \leq \dim(\Omega_R^i(M))$, while on the other hand $\dim(\Omega_R^i(M)) \leq \dim(R)$, so our result improves the upper and lower bounds for the degree of $\tau_i^{M,R,\mathfrak{m}}(n)$ given in [3]. Furthermore, if M has a rank and $N = R$, it

is shown in [5], that $\deg(\tau_i^{M,R,\mathfrak{m}}(n)) = \dim(R) - 1$. Since in this case, R , M and $\Omega_R^i(M)$ all have the same dimension, Theorem 3.12 recovers this result as well.

4. MORE GENERAL FILTRATIONS

We now turn to giving an analogue of the main results in [5] for $\text{Ext}^i(M, N/I^n N)$. In [5] the second and fourth authors considered the Hilbert polynomial giving the lengths of $\text{Tor}_i(M, R/I^n)$. In that paper, various assumptions were made on I and M which forced $\tau_i^{M,N,I}(n)$ to have maximal degree $\ell(I) - 1$. Roughly speaking the assumptions on M were made so that the i^{th} syzygy has maximal dimension. The assumptions on the filtrations given in [5] were made in order to replicate some of the properties satisfied by the \mathfrak{m} -adic filtration. The reason for this is now clear in light of Theorem 3.12. Likewise, we may use some of the ideas underlying Theorem 3.2 to give the corresponding results for $\mathcal{E}_{M,N,I}^i(n)$, for similar I and M .

Before presenting our main results in this section, we state a proposition which is simply a restatement of Proposition 2.1 in the the context of the contravariant extension functor. For the sake of consistent notation with the previous section, we set $T_i := \text{im}(\overline{\phi_{i+1}})$ and

$$\mathcal{T}_i := \bigoplus_{n \geq 0} (I^n N^{\beta_{i+1}(M)} \cap T_i) / I^n T_i.$$

Note that in our present context, \mathcal{T}_i is just \mathcal{M} from Proposition 2.1.

Proposition 4.1. *Let (R, \mathfrak{m}) be a local ring and suppose that M and N are finitely generated R -modules. Fix $0 \leq i \leq \text{p.d.}(M)$ and let $I \subseteq R$ be an ideal. Assume that for all large n , the lengths of $\text{Ext}^i(M, N/I^n N)$ are finite. Then*

$$\deg(\mathcal{E}_{M,N,I}^i(n)) = \max\{\dim(\text{Ext}^i(M, N)), \dim(\mathcal{T}_i) - 1\}.$$

We now give an analogue of Theorem 3.2 for ideals divisible by \mathfrak{m} .

Theorem 4.2. *Let (R, \mathfrak{m}) be a local ring and suppose that M and N are finitely generated R -modules. Let I be an ideal divisible by \mathfrak{m} , i.e., $I = \mathfrak{m}C$, for some ideal $C \subseteq R$. Fix $0 \leq i \leq \text{p.d.}(M)$ and suppose that the lengths of $\text{Ext}^i(M, N/I^n N)$ are finite for n large. Then*

$$\deg(\mathcal{E}_{M,N,I}^i(n)) = \max\{\dim(\text{Ext}^i(M, N)), \ell_{T_i}(I) - 1\},$$

where as before, $T_i := \text{im}(\overline{\phi_{i+1}})$.

Proof. By Proposition 4.1, it suffices to prove $\dim(\mathcal{T}_i) = \ell_{T_i}(I)$. Consider the filtration \mathcal{J} whose terms are $J_n := \mathfrak{m}^{n-1}C^n$, $n \geq 1$. Note that $J_{n+1} = IJ_n$ for all $n \geq 1$. Then \mathcal{J} is an I -good filtration, so by Proposition 2.2 in [5], the graded module $\bigoplus_{n \geq 0} J_n T_i / \mathfrak{m} J_n T_i$ has dimension $\ell_{T_i}(I)$. Now, on the one hand, since \mathbf{F} is a minimal resolution, $J_n T_i \subseteq I^n N^{\beta_{i+1}(M)} \cap T_i$. Thus,

$$\bigoplus_{n \geq 0} J_n T_i / \mathfrak{m} J_n T_i \subseteq \mathcal{T}_i,$$

from which it follows that \mathcal{T}_i has dimension at least $\ell_{T_i}(I)$. On the other hand, set $S := R / \text{ann}(T_i)$. Then \mathcal{T}_i is a finitely generated module over the Rees algebra $\mathcal{R}(IS)$ of S with respect to IS whose graded components have finite length. Thus, there exists $r > 0$ such that $\mathfrak{m}^r \mathcal{R}(IS)$ annihilates \mathcal{T}_i . Therefore,

$$\dim(\mathcal{T}_i) \leq \dim(\mathcal{R}(IS) / \mathfrak{m}^r \mathcal{R}(IS)) = \ell_S(I) = \ell_{T_i}(I).$$

Thus, $\dim(T_i) = \ell_{T_i}(I)$, which gives what we want. \square

Remark 4.3. For $I = \mathfrak{m}C$, the value of $\ell_{T_i}(I)$ can vary anywhere between zero and $\dim(T_i)$ and the latter can be as large as $\dim(R)$. However, if we set $S := R/\text{ann}(T_i)$ and assume that $\text{height}(IS) > 0$, then $\ell_{T_i}(I)$ achieves its maximum value of $\dim(T_i)$. To see this, after a change in notation, it suffices to see that if $\text{height}(I) > 0$, then $\ell(I) = d = \dim(R)$. For this, recall that since $\text{height}(C) > 0$, the Hilbert-Samuel polynomial giving the lengths of the modules $C^r \mathfrak{m}^s / C^r \mathfrak{m}^{s+1}$ for r, s large is a polynomial of total degree $d - 1$ in r and s with non-negative leading coefficients. For large n , we set $n := r = s$. It follows that the polynomial giving the lengths of $(C\mathfrak{m})^n / \mathfrak{m}(C\mathfrak{m})^n = I^n / \mathfrak{m}I^n$ has degree $d - 1$. Therefore, $\ell(I) = d$.

If $N = R$, we have an immediate corollary for ideals divisible by \mathfrak{m} .

Corollary 4.4. *Let R be a local ring and M a finitely generated R -module. Assume that R is unmixed and equidimensional. Assume further that either M has a rank or R is generically Gorenstein. Let $I = \mathfrak{m}C$ be an ideal of R such that $\text{height}(I) > 0$. Then, $\deg(\mathcal{E}_{M,R,I}^i(n)) = d - 1$.*

Proof. If M has a rank or R is generically Gorenstein, $\dim(\text{Ext}^i(M, R)) \leq d - 1$. Thus, by Theorem 4.2, it suffices to show that $\ell_{T_i}(I) = d$. Consider the $(i + 1)^{\text{st}}$ syzygy of M , $K_i := \text{im}(\phi_{i+1})$. Since K_i and T_i have the same support, we just have to show that $\ell_{K_i}(I) = d$. Let P be any prime minimal in the support of K_i . Since R is unmixed and equi-dimensional, $S := R/P$ has dimension d . Since $\text{height}(I) > 0$, $\text{height}(IS) > 0$. By the remark above, $\ell_S(I) = d$. Since $\ell_{K_i}(I)$ is the maximum value over all such S , it follows that $\ell_{K_i}(I) = d$, which is what we want. \square

Our final goal is to state a theorem that is a variant for the contravariant extension functor of the main results in [5]. It gives a number of cases where the degree of $\mathcal{E}_{M,N,I}^i(n)$ is $d - 1$. First, we require a definition and a lemma. In the lemma we maintain the notation established throughout the paper.

Definition 4.5. Let M be a finitely generated R -module. M is said to *test finite projective dimension* if for all finitely generated modules N , N has finite projective dimension if and only if for some $i > 0$, $\text{Tor}_i^R(M, N) = 0$.

While the residue field \mathfrak{k} obviously satisfies this condition - and this is the case one is often interested in - it follows from [1] Corollary 3.3, that R/J satisfies the condition for any integrally closed \mathfrak{m} -primary ideal J . Of course, if M tests finite projective dimension, then so does any syzygy of M .

Lemma 4.6. *Let N be a finitely generated R -module such that $N_P \neq 0$, for every minimal prime P . Let M be a finitely generated R -module such that either M has a rank or M is free of constant rank on the punctured spectrum of R and M tests finite projective dimension. Assume further that $i < \text{p.d.}(M)$. Then for $\overline{\phi_{i+1}}$ as in section three and $T_i := \text{im}(\overline{\phi_{i+1}})$, the annihilator of T_i is nilpotent.*

Proof. Set $K_i := \text{im}(\phi_{i+1})$. Then $(K_i)_P \neq 0$ for all minimal primes $P \subseteq R$. When M has a rank, this follows from Remark 2.1 in [5]. If M is free of constant rank on the punctured spectrum of R and M tests finite projective dimension, this was shown for finite length M in the first paragraph of the proof of Theorem 3.3 in [5], but for the reader's convenience, we repeat the argument in this slightly more general case. First, note that if $\text{depth}(R) > 0$, then M has a rank, and we are in the

previous case. Suppose $\text{depth}(R) = 0$. By hypothesis, K_i is also free of constant rank on the punctured spectrum of R . If this locally constant rank were zero, then K_i would have finite length. But then by Corollary 2.3 (c), $\text{Tor}_i(R/\mathfrak{m}^n, M) = 0$ for large n . By the hypothesis on M , R/\mathfrak{m}^n has finite projective dimension, which cannot be when $\text{depth}(R) = 0$. Thus K_i does not have finite length, and thus must be nonzero when localized at any non-maximal prime ideal. In particular, $(K_i)_P$ is non-zero for each minimal prime P . Thus, in both cases, the map $(\phi_{i+1})_P$ is non-zero for each such P and so its transpose $(\phi_{i+1}^t)_P$ is also non-zero. Since the \mathbf{F}_P is split exact for all minimal primes P , the complex $\text{Hom}(\mathbf{F}, N)_P$ is also split exact, and since N_P is non-zero for each minimal prime P , it follows that $(T_i)_P$ is not zero for each minimal prime P . Thus, the annihilator of T_i is nilpotent. \square

Theorem 4.7. *Let (R, \mathfrak{m}) be a local ring of dimension d and $I \subseteq R$ an ideal having analytic spread d . Let N and M be finitely generated R -modules such that $\lambda(\text{Ext}^i(M, N/I^n N))$ is finite for n large. Here, $0 < i < \text{p.d.}(M)$. Assume that M has a rank (possibly zero) or M has is free of constant rank on the punctured spectrum of R and M tests finite projective dimension. Assume further that N_P is non-zero for every minimal prime P . Suppose that one of the following conditions is satisfied.*

- (i) $I = \mathfrak{m}C$ for some ideal $C \subseteq R$.
- (ii) $(\mathfrak{m}I^n N :_N \mathfrak{m}) = I^n N$, for large n .
- (iii) $(I^n :_R \mathfrak{m}) \not\subseteq \overline{I^n}$, for some n and R is quasi-unmixed.

Then $\deg(\mathcal{E}_{M,N,I}^i(n)) = d - 1$.

Proof. We first note that either assumption on M yields $\dim(\text{Ext}^i(M, N)) \leq d - 1$. Thus, by Proposition 4.1, we must show that $\dim(\mathcal{T}_i) = d$. The proof of this for each of the stated conditions follows closely the proofs given for Theorems 3.3 and 3.4 in [5]. We will try to give a convincing account without repeating all of the details from [5]. A crucial point in each case is that the annihilator of T is nilpotent, by Lemma 4.6.

Now suppose $I = \mathfrak{m}C$, for some ideal C . From the proof of Theorem 4.2, we know that $\dim(\mathcal{T}_i) = \ell_{T_i}(I)$. By Lemma 4.6, T has nilpotent annihilator. Thus $\ell_{T_i}(I) = \ell(I) = d$, which gives what we want.

Suppose $(\mathfrak{m}I^n N :_N \mathfrak{m}) = I^n N$ for large n . Replacing I by I^t for t sufficiently large allows us to show, just as in paragraph three in the proof of [5], Theorem 3.3, that for all n , we have an equality of socles,

$$\text{Soc}(\mathcal{T}_i) = \text{Soc}(I^{n-1}T_i/I^n T_i).$$

Since T_i has nilpotent annihilator, $d = \ell(I) = \ell_{T_i}(I)$, and the same proof used in paragraph four on page 3079 in [5] shows that the module $\bigoplus_{n \geq 0} \text{Soc}(I^{n-1}T_i/I^n T_i)$ has dimension d . This in turn implies that \mathcal{T}_i also has dimension d , which is what we want.

Finally, suppose that $(I^n : \mathfrak{m}) \not\subseteq \overline{I^n}$ for some n and that R is quasi-unmixed. Again, since the resolution \mathcal{F} is a minimal resolution, $(I^n : \mathfrak{m})T_i \subseteq I^n N^u \cap T_i$. Thus,

$$\bigoplus_{n \geq 0} (I^n : \mathfrak{m})T_i/I^n T_i \subseteq \mathcal{T}_i.$$

Since the annihilator of T_i is nilpotent, the same proof used in paragraphs three and four in the proof of [5], Theorem 3.4, shows that the module $\bigoplus_{n \geq 0} (I^n : \mathfrak{m})T_i / I^n T_i$ has dimension d . Therefore, \mathcal{T}_i also has dimension d , and the proof is complete. \square

Remark 4.8. One should note that no assumption about the non-vanishing of the extension modules $\text{Ext}^i(M, N/I^n N)$ in the theorem is made. Thus, in particular, the theorem shows that for ideals and modules as in the theorem, $\text{Ext}^i(M, N/I^n N)$ is not zero. A similar remark applies to the corresponding torsion modules.

Remark 4.9. Since N is not injective, one does not automatically obtain the analogue of results for $\text{Ext}^{i+1}(M, I^n N)$ from $\text{Ext}^i(M, N/I^n N)$ by dimension shifting. However, for the question we're interested in, if we assume that the modules $\text{Ext}^i(M, N)$, $\text{Ext}^i(M, N/I^n N)$, and $\text{Ext}^{i+1}(M, I^n N)$ all have finite length for large n , the answer follows readily. Indeed, for all $n \geq 1$, the short exact sequence

$$0 \rightarrow I^n N \rightarrow N \rightarrow N/I^n N \rightarrow 0$$

gives rise to the long exact sequence

$$\text{Ext}^i(M, N) \rightarrow \text{Ext}^i(M, N/I^n N) \rightarrow \text{Ext}^{i+1}(M, I^n N) \rightarrow \text{Ext}^{i+1}(M, N).$$

It quickly follows that the polynomials giving the lengths of $\text{Ext}^i(M, N/I^n N)$ and $\text{Ext}^{i+1}(M, I^n N)$ differ only by a constant and consequently have the same degree, assuming both lengths are not zero. We use this for the corollaries below.

In [6], page 763, it is asked whether, for fixed i , the polynomial giving the i^{th} Betti number of $I^n N$ has degree $\ell_N(I) - 1$, provided it is not identically zero. One of the purposes of [5] was to show that the answer is yes in a number of cases. Similarly, by taking $M = \mathfrak{k}$ in Theorem 4.7, we can now note that for fixed i the polynomial giving the i^{th} Bass number of $I^n N$ is given by a polynomial of degree $d - 1 = \ell_N(I) - 1$ in essentially the same cases.

Corollary 4.10. *Let (R, \mathfrak{m}) be a local ring of dimension d and N a finitely generated R -module such that $N_P \neq 0$, for all minimal primes P . Assume $I \subseteq R$ satisfies $\ell(I) = d$ and any one of the conditions (i)-(iii) from Theorem 4.7. Then for any $i > 0$ with i less than the projective dimension of \mathfrak{k} , the i^{th} Bass numbers of $N/I^n N$ and $I^n N$ are given by polynomials of degree $d - 1$. \square*

The following instance of the previous corollary deserves special attention.

Corollary 4.11. *Let (R, \mathfrak{m}) be a quasi-unmixed local ring and I an integrally closed \mathfrak{m} -primary ideal. Then for any $i > 0$ with i less than the projective dimension of \mathfrak{k} , the i^{th} Bass numbers of R/I^n and I^n are given by polynomials of degree $d - 1$. \square*

5. APPENDIX

As mentioned in section three above, our results concerning the degree of $\mathcal{E}_{M, N, \mathfrak{m}}^i(n)$ are closely related to the dimension of the corresponding syzygies associated with M . As a consequence, we can shed some light on the following interesting question.

Question 5.1. Let (R, \mathfrak{m}) be a local ring and M a finitely generated R -module. Is the dimension of the n^{th} syzygy $\Omega_R^n(M)$ stable for n sufficiently large?

In the following remark, we record a few easy observations concerning Question 5.1.

Remark 5.2. (i) If R is unmixed and equidimensional, then clearly all syzygies have dimension equal to $\dim(R)$.

(ii) If the dimension of syzygies is ultimately constant, then that constant value must equal $\dim(R)$. This follows since given two consecutive syzygies, one of them must have dimension equal to $\dim(R)$.

(iii) If $\beta_i(M) > \beta_{i-1}(M)$, then $\dim(\Omega_R^{i+1}(M)) = \dim(R)$. This is because for any prime ideal P , $(\phi_i)_P$ cannot be injective. Thus, in fact, $\Omega_R^{i+1}(M)_P \neq 0$, for all P , so in this case $\Omega^{i+1}(M)$ has nilpotent annihilator. It follows that if the betti numbers of M are eventually increasing, then Question 5.1 has a positive answer for M .

(iv) If $\beta_i(M) < \beta_{i-1}(M)$, then $\dim(\Omega^{i-1}(M)) = \dim(R)$. This is because for any prime ideal P , $(\phi_i)_P$ cannot be surjective. Thus, in fact, $\Omega^{i-1}(M)_P \neq 0$, for all P , so in this case, $\Omega^{i-1}(M)$ has nilpotent annihilator.

Proposition 5.3. *Let (R, \mathfrak{m}) be a local ring of dimension d and M a finitely generated R -module. Let $\Omega_R^{i+1}(M)$ be an $(i+1)^{st}$ syzygy of M with $1 \leq i < \text{p.d.}(M)$. Assume that $\beta_i(M) = \beta_{i-1}(M)$. Then*

$$\max\{\dim(\text{Ext}^i(M, R)), \dim(\Omega_R^{i+1}(M)) - 1\} \geq d - 2.$$

In particular, if $\dim(R) \geq 2$ and M is free of constant rank on the punctured spectrum of R , then in fact, $\dim(\Omega_R^{i+1}(M)) = d$.

Proof. We may assume that $\Omega_R^{i+1}(M) = \text{im}(\phi_{i+1})$. By Theorem 3.2

$$\deg(\mathcal{E}_{M,R,\mathfrak{m}}^i(n)) = \max\{\dim(\text{Ext}^i(M, R)), \dim(T_i) - 1\},$$

where T_i now denotes the image of the transpose of ϕ_{i+1} in $R^{\beta_{i+1}(M)}$. On the other hand, $\dim(T_i) = \dim(\Omega_R^{i+1}(M))$, since for any prime P the matrix $(\phi_{i+1})_P$ is the zero matrix if and only if its transpose is the zero matrix. Therefore,

$$\deg(\mathcal{E}_{M,R,\mathfrak{m}}^i(n)) = \max\{\dim(\text{Ext}^i(M, R)), \dim(\Omega_R^{i+1}(M)) - 1\}.$$

The first statement in the corollary now follows from Corollaries 3.5(c) and 3.7.

To prove the second statement, we note that $\dim(\text{Ext}^i(M, R)) = 0$ since M is free on the punctured spectrum. Therefore, if $d \geq 3$, the maximum value above is $\dim(\Omega_R^{i+1}(M)) - 1$. If $d = 2$, the maximum value is again $\dim(\Omega_R^{i+1}(M)) - 1$. For this, it is enough to show that the dimension of $\Omega_R^{i+1}(M)$ is positive. Assume by way of contradiction, that $\dim(\Omega_R^{i+1}(M)) = 0$. Consider the following exact sequence

$$0 \rightarrow \Omega_R^{i+1}(M) \rightarrow R^{\beta_i(M)} \xrightarrow{\phi_i} R^{\beta_{i-1}(M)} \rightarrow \Omega_R^{i-1}(M) \rightarrow 0,$$

where $\beta_i(M) = \beta_{i-1}(M)$. Since $\Omega_R^{i+1}(M)$ has finite length and $\Omega_R^{i-1}(M)$ is free of constant rank on the punctured spectrum, it follows from the sequence above that $\Omega_R^{i-1}(M)$ also has finite length. Thus, the $\det(\phi_i)$ generates an \mathfrak{m} -primary ideal, so $\dim(R) \leq 1$, which is a contradiction. Therefore $\dim(\Omega_R^{i+1}(M))$ is strictly positive and hence

$$\dim(\Omega_R^{i+1}(M)) - 1 = \max\{\dim(\text{Ext}^i(M, R)), \dim(\Omega_R^{i+1}(M)) - 1\} \geq d - 2,$$

where the inequality follows from the first statement. Thus, $\dim(\Omega_R^{i+1}(M)) \geq d - 1$. To improve this, note that for any prime ideal $P \neq \mathfrak{m}$, $\Omega_R^{i+1}(M)_P$ is a free R_P -module of rank independent of P . If this rank were zero, then $\Omega_R^{i+1}(M)$ would have finite length. But since $\dim(\Omega_R^{i+1}(M)) \geq 1$, this cannot be. Thus, the constant

rank of each $\Omega_R^{i+1}(M)_P$ is not zero, so $\Omega_R^{i+1}(M)_P$ is not zero for all non-maximal primes P . In particular, $\dim(\Omega_R^{i+1}(M)) = d$. \square

The following example shown to us by Hamid Rahmati shows that we cannot relax the hypothesis $\dim(R) \geq 2$ in the second statement of Corollary 5.3.

Example 5.4. Let $R := k[[x, y]]/(x^2, xy)$. Let $M := R/(y)$, so that M is a finite length R -module. Consider the start of a free resolution of the R -module M

$$0 \rightarrow xR \rightarrow R \xrightarrow{y} R \rightarrow M \rightarrow 0.$$

Since x is a socle element, the second syzygy of M also has finite length, and so $\dim(\Omega_R^2(M)) < \dim(R)$.

Proposition 5.5. *Let (R, \mathfrak{m}) be a local ring and M a finitely generated R -module.*

- (a) *If M has non-decreasing betti numbers, then M has at most one syzygy $\Omega_R^j(M)$ with finite length. Moreover, $1 \leq j \leq d$.*
- (b) *If the betti numbers of M are eventually non-decreasing, then M has only finitely many syzygies with finite length.*

Proof. For (a), suppose that M has non-decreasing betti numbers and $M' := \Omega_R^j(M)$ is a syzygy of finite length. Then M' is free of constant rank zero on the punctured spectrum of R , so that by Proposition 5.3, $\dim(\Omega_R^{i+1}(M')) = d$, for all $i \geq 1$. Thus, $\dim(\Omega_R^s(M)) = d$, for all $s \geq j + 2$. On the other hand, the exact sequence

$$0 \rightarrow \Omega_R^{j+1}(M) \rightarrow F_j \rightarrow \Omega_R^j(M) \rightarrow 0$$

shows that $\dim(\Omega_R^{j+1}(M)) = d$, so that $\dim(\Omega_R^s(M)) = d$, for all $s \geq j + 1$. Note that this argument now precludes the possibility of $\Omega_R^i(M)$ having finite length for some $i < j$, so that M has at most one syzygy of finite length. To see that $j \leq d$, suppose to the contrary that $j > d$. Since $\Omega_R^j(M)_P = 0$, for all prime ideals $P \neq \mathfrak{m}$, $\text{p.d.}(M_P) < \infty$, for all primes $P \neq \mathfrak{m}$. It follows from this that $\Omega_R^{d-1}(M)$ is free on the punctured spectrum of R . Note that since $\Omega_R^j(M)$ has finite length and $j > d$, working backwards from $\Omega_R^j(M)$ it follows that $\Omega_R^{d-1}(M)$ must also have constant rank on the punctured spectrum. If we now apply Corollary 5.3 to $\Omega_R^{d-1}(M)$, it follows that $\dim(\Omega_R^j(M)) = d$, and this is a contradiction. Thus, $j \leq d$, as required. This finishes the proof of part (a) of the proposition.

Finally, part (b) of the proposition follows immediately from the proof of the first statement in part (a). \square

Remark 5.6. It is clear from Proposition 5.5 that Question 5.1 is related to another more important question, namely whether or not every finitely generated module over an arbitrary local ring has the property that its betti numbers are eventually non-decreasing. Suppose this latter property were true for all local rings. Replacing the module M by one of its large syzygies, one could assume that the betti numbers of M were non-decreasing. If $\Omega_R^i(M)$ were a syzygy of dimension less than d , localize at prime P minimal in its support. By Corollary 5.3 and Proposition 5.5, $\dim(\Omega_R^s(M)_P) = \dim(R_P)$, for all $s \geq i + 1$. Thus, if R were equidimensional, then $\Omega_R^s(M)$ would have dimension d , for all $s \geq i + 1$, and the dimension of the syzygies of M would stabilize.

REFERENCES

1. A. Corso, C. Huneke, D. Katz and W. Vasconcelos *Integral closure of ideals and annihilators of homology*, Lecture Notes in Pure and Applied Mathematics **227** (2005), 33-48.
2. A. Crabbe and J. Striuli, *Constructing big indecomposable modules*, Proc. Amer. Math. Soc., to appear.
3. S. Iyengar and T. Puthenpurakal, *Hilbert-Samuel functions of modules over Cohen-Macaulay local rings*, Proc. Amer. Math. Soc. **135**, No. 4 (2007), 637-648.
4. S. Goto, F. Hayasaka and R. Takahashi, *On vanishing of certain Ext modules*, JI. Math. Soc. Japan **60**, No. 4 (2008), 1045-1064.
5. D. Katz and E. Theodorescu, *On the degree of Hilbert polynomials associated to the torsion functor*, Proc. Amer. Math. Soc. **135**, No. 10 (2007), 3073-3082.
6. V. Kodiyalam, *Homological invariants of powers of an ideal*, Proc. Amer. Math. Soc. **118**, No. 3 (1993), 757-764.
7. E. Theodorescu, *Derived functors and Hilbert polynomials*, Math. Proc. Camb. Phil. Soc. **132** (2002), 75-88.

DEPARTMENT OF MATHEMATICS, SYRACUSE UNIVERSITY, SYRACUSE, NY 13244-1150
E-mail address: `amcrabbe@syr.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, LAWRENCE, KS 66045
E-mail address: `dlk@math.ku.edu`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, FAIRFIELD UNIVERSITY, FAIRFIELD,
CT 06824
E-mail address: `jstriuli@mail.fairfield.edu`

DIVISION OF STATISTICS, NORTHERN ILLINOIS UNIVERSITY, DE KALB, IL 60115
E-mail address: `theodore@math.niu.edu`